



72 Lowry Street, Rochester, Vic, 3561
Ph: (03) 54842255 Fax: (03) 54842424 Email: admin@rochyvet.com.au

Vaccination Protocols

DOGS

The C3 vaccination protects dogs against canine parvovirus, distemper virus and hepatitis.

The BBPi and KC vaccinations protect dogs against canine cough, which is a requirement for dogs going into kennels. Using these vaccinations in combination with the C3 vaccine is termed a C5 vaccination schedule.

Puppies - 1st vaccination with C3 at 6-8 weeks +/- BBPi
2nd vaccination with C3 at 10-12 weeks +/- BBPi

For pups that are 10 weeks or more that require canine cough vaccination

- 1st Vaccination Nobivac DHP plus BBPi
- 2nd Vaccination Nobivac BBPi only 4 weeks later

Alternatively they can have just C3 plus KC Intranasal vaccination.

At 15mths they will receive a C3 with or without Nobivac BBPi or Nobivac KC intranasal

Adults – Annual vaccination with either single KC intranasal vaccination or if they have had the 1ry course of two BBPi injections then can give BBPi

Tri-annual booster of C3 with or without Canine cough booster

CATS

The F4 vaccination protects cats against feline calici virus, feline rhinotracheitis (feline herpes virus), panleucopenia and *feline chlamydia psittaci*.

Vaccination schedule:

Kittens - 1st vaccination at 8 weeks or older with F4
2nd vaccination 4 weeks later with F4

Adults - Annual vaccination with F4

All previously unvaccinated cats require 2 doses of F4 given 4 weeks apart for full protection against cat flu

For vaccination against Feline Immunodeficiency Virus (FIV)

Kittens - 1st vaccination at 8 weeks with F4 + FIV
2nd vaccination 10 weeks with FIV
3rd vaccination 12 weeks with F4 + FIV

Adults - Annual vaccination with F4 + FIV

RABBITS

Cylap vaccine protects rabbits against haemorrhagic viral disease (calicivirus)

First vaccination at 10-12 wks, then annual vaccination

If first vaccination is given prior to 10 weeks of age then give 2nd injection 4 weeks later and repeat annually